

ИНСТИТУТ ЯДЕРНОЙ ФИЗИКИ СО АН СССР

D.L.Shepelyansky

QUANTUM DIFFUSION LIMITATION
AT EXCITATION OF RYDBERG ATOM IN
VARIABLE FIELD

PREPRINT 83-61



НОВОСИБИРСК

INSTITUTE OF NUCLEAR PHYSICS

Siberian Division

USSR ACADEMY OF SCIENCES

Preprint 83-61

D.L.Shepelyansky

QUANTUM DIFFUSION LIMITATION AT EXCITATION
OF RYDBERG ATOM IN VARIABLE FIELD

Novosibirsk

1983

QUANTUM DIFFUSION LIMITATION AT EXCITATION
OF RYDBERG ATOM IN VARIABLE FIELD

D.L. Shepelyansky

Institute of Nuclear Physics
630090, Novosibirsk 90, USSR

Abstract

A computer simulation method is used to investigate excitation of a hydrogenlike atom in the field of an electromagnetic wave from the states with $n \sim 50$ and parabolic quantum numbers $n_1 \gg n_2$ (or $n_1 \ll n_2$). It is shown that diffusion over levels is much slower than in the classical case. A significant part in atom excitation is played by multiphoton resonances with the number of photons $k \gg 10$.

I. Introduction

For the last few years a number of interesting experiments was carried out on ionization and excitation of hydrogen atoms by the microwave field from the states with the main quantum number $n \sim 50$ /1-3/. The first of these experiments showed at the frequency $\omega/2\pi = 9.9$ GHz a strong ionization of atoms with $n \approx 65$ at the electric field $\mathcal{E} \approx 0.06n^4$ and adiabatic tunneling parameter $\gamma = \omega/\mathcal{E}n \approx 7/4$ ($\omega = 0.43n^3$)*. To explain the results of the experiment a diffusion mechanism of atom ionization was offered in /5/. The cause of electron diffusion arising in an atom due to the purely monochromatic field effect can be understood in the following way. It is quite possible to assume /6/ that at $n \approx 50 \gg 1$ the quasiclassical approximation is good enough and to pass on to investigation of the classical system. The latter is substantially nonlinear and at field $\mathcal{E} \gg \mathcal{E}_c$ there arises a stochasticity leading to a diffusion excitation of an electron

* Here and below we use atomic units; n, l, m are principal, orbital and magnetic quantum numbers.

/7/. In case of linear and circular polarizations and $\omega \approx \mathcal{E} = n^3$ the estimation for \mathcal{E}_c was obtained in /7,8/ on the grounds of the overlap criterium /9/. Dependences of critical field and diffusion rate on frequency are given in /10,11/. Despite all attractiveness of this classical approach, its only justification is the coincidence of the relative portion of ionized atoms at $\mathcal{E}n \approx 0.06$ /1/ with the probability of ionization obtained through computer simulation of classical system's dynamics /6/. The accuracy of coincidence is about 30%.

At the same time a number of numerical experiments with simple quantum systems /12,13/ have shown that statistical properties of quantum dynamics are much weaker than those of classical one. Moreover in the course of time the effect of quantum corrections increases /14/ and leads to diffusion's slowing down and its final almost complete stopping. First this effect of quantum diffusion limitation was observed in the model of quantum rotator /12/. The same phenomenon can take place at diffusion ionization of atom with $n \gg 1$. Preliminary estimates /11/ show that relative magnitude of the quantum corrections can be compared to unity and so there may be a significant distinction between quantum dynamics of atom excitation and classical one.

II. The Model of Surface-State Electrons

When investigating quantum and classical dynamics we employed the computer simulation method. We chose an external field linear polarized along the axis z . The atom was initially excited into states with quantum numbers which satisfied the following conditions $n \sim 50$, $m=0$, $n_1 \gg n_2$ (or $1 \ll l \ll n^2$). For these states the ratio of cross-size to longitudinal one is small and that is why it's possible to describe the dynamics of excitation in this case by the one-dimensional Hamiltonian

$$H = \frac{p^2}{2} - \frac{1}{|z|} + \mathcal{E} z \cos \omega t \quad (1)$$

with the boundary condition $\psi(0)=0$, which corresponds to a infinite potential wall at $z=0$. This Hamiltonian also describes the excitation of states of atom with $l=0$ in the external field with the potential $V = \mathcal{E}|z| \cos \omega t$. Consequently for (1) matrix elements $Z_{nn'} = Z_{nn'}$. To prove the fact that such a modification of the initial Hamiltonian won't lead to any significant change of the dynamics we can present the following arguments:

1. When $1 \ll l \ll n^2$ the matrix elements $Z_{nn'}^{\mathcal{E}z}$ don't depend on l /15/, as a result of which their sum Z_Z that determines the probability of transition from one shell into another in dipole approximation to within several

per cent is equal to $r_{nn'}$ ($n, n' \sim 50$). So, for $n=40, n'=41, l=2$ the ratio $z_x/r_{nn'} = -0.996$ (here we use for z_x its quasiclassical value).

2. Due to Coulomb degeneration the transition frequency between adjoining sublevels of one and the same shell prove much smaller than Kepler frequency: $3\epsilon n/2\Omega \ll 0.1 \ll \ll 1$ for $\epsilon_n = \epsilon n^2 \leq 0.06$. So, despite the fact that the width of Stark multiplet exceeds the distance between the shells $\Delta\Omega \gg \Omega$ the motion of sublevels proves synchronized and it's possible to substitute the whole shell for one level. A proof of such approximation is given by the experiments /2,3/, where the observed resonant picture of excitation in the field with $\epsilon_n \approx 0.06$ corresponded to the unperturbed spectrum of atom levels.

3. In the classical system when $l \ll n$ the dependence of ϵ_{cr} and the rate of diffusion on l proves insignificant /11/. The Hamiltonian (1) describes precisely the motion for the orbits with eccentricity $e=1$ stretched along the field. In the quantum system the states with parabolic quantum numbers $n_1=n-1, n_2=0$ (or vice versa) and close to them ($n_1 \gg n_2$) correspond to such classical orbits. Let's see if this condition will be violated in the course of time. To do this we shall consider unperturbed ($\epsilon=0$) phase-action variables corresponding to parabolic quantum numbers. For these variables

$$Z = \frac{3}{2} n(n_1 - n_2) + \sum_{\substack{m_1, m_2 \\ m_1 + m_2 \neq 0}} B_{m_1 m_2} \exp[i(m_1 \lambda_1 + m_2 \lambda_2)]$$

$$B_{m_1 m_2} = \frac{1}{m_1 + m_2} \left[\mu_2 J_{m_1}(\mu_1(m_1 + m_2)) J'_{m_2}(\mu_2(m_1 + m_2)) - \right. \quad (2)$$

$$\left. - \mu_1 J'_{m_1}(\mu_1(m_1 + m_2)) J_{m_2}(\mu_2(m_1 + m_2)) \right],$$

$\mu_{1,2} = [n_{1,2}(n - n_{2,1})/n^2]^{1/2}$; λ_1, λ_2 - phases conjugated to n_1, n_2 ; where the derivative is taken over the argument of Bessel function. According to the correspondence principle the values $B_{m_1 m_2}$ determine matrix elements $z_{n_1 n_2}$ of the transition between the unperturbed levels with $n_{1,2} = n_{1,2} - n_{1,2}$. When $n_1 \gg n_2$ ($m=0$) matrix elements for the transitions with the change of n_2 contain a supplementary small parameter n_2/n . The change magnitude of n_2 can be estimated in the course of time in the following way. For $\omega \approx \approx \Omega$ and $\epsilon > \epsilon_{cr}$ there starts a diffusion electron excita-

tion. According to the theoretical estimates /11/ the rate of diffusion over n and the value of ϵ_{cr} are:

$$D_n = \frac{d(\Delta n)^2}{dt} \approx \frac{1}{3} \epsilon^2 n^2; \quad \epsilon_{cr} \approx \frac{1}{84 n^4} \quad (3)$$

When $\epsilon > \epsilon_{cr}$ the phase λ_1 changes randomly in the course of time, which makes it possible to estimate the rate of diffusion over n_2 : $D_{n_2} \sim (n_2/n)^2 D_n$. From the above we see that in the course of a numerical experiment $\tau = \omega t/2\pi \leq 50$ at $\epsilon n^2 \approx 0.04, n_2 \sim 1$ the change $\Delta n_2 \sim 1$ and so the one-dimensional approximation is not violated. It's interesting to note that when $\epsilon n^2 n_2 \ll 1$ the dynamics over the second degree of freedom (n_2, λ_2) proves substantially quantum, since in this case the perturbation theory can be used /16/: $B_{m_1 m_2}/\Omega \ll 1$ ($m_2 \neq 0$). Let's point out the fact that when $m_1 + m_2 = \text{const} \sim 1$ the matrix elements $B_{m_1 m_2}$ with $m_1, m_2 \gg 1$ are exponentially small. This proves still more rigorously the effect observed in the experiments where the shell shows itself as one level /2,3/.

To determine in quantitative terms how good is the selected one-dimension approximation we made a comparison with the results of numerical experiments for a real atom in a linear polarized field carried out by P.M. Izrailev. The initial condition was given by parabolic quantum numbers $n_1 = n_0 - 1, n_2 = 0$. The comparison was made for $n_0 = 10, 9$ and showed a good agreement with the results obtained in a one-dimensional approximation. So, for instance, the probabilities of excitation $W_{n_1 > n_0, n_2}$ into states with $n > n_0$ coincide with the accuracy of $\approx 10-20\%$.

Hence, proceeding from the above arguments we can conclude that the system (1) will describe in qualitative and quantitative terms the dynamics of Rydberg atom excitation from states $n_1 \gg n_2$ or $n_1 \ll n_2$ (and qualitatively from $1 \leq l \ll n^{1/2}$). Besides, this model describes exactly the dynamics of surface-state electrons excitation /17/ in a variable field, directed perpendicular to the surface of liquid helium and that is why its investigation is of interest in itself. It was offered to carry out an experiment with such a system in reference /18/*.

III. Numerical Experiments

Simulation of quantum dynamics of system (1) was carried out in the following way. At the initial moment of time one level of the unperturbed system with $n_0 \sim 50$ was

*) We note that the functional dependence $\epsilon_{cr}(\omega)$ in /18/ was not found correctly (there is a mistake in f.(8)). See for a correct answer /11/.

excited. The field frequency was close to the Kepler one: $\omega n_0^2 = \omega_0 \sim 1$ and the field $\epsilon n_0^2 = \epsilon_0 \sim 0.06$. Further on we carried out a numerical integration of equations for amplitudes C_n of the unperturbed Hamiltonian's states.

$$i \dot{C}_n = -\frac{1}{2n^2} C_n + \sum_{n'=n_{\min}}^{n_{\max}} V_{nn'} C_{n'} \quad (4)$$

Thus, levels with $n_{\min} \leq n \leq n_{\max}$ only were involved in the dynamics. Since at the cited values of parameters ϵ and ω diffusion in the classical system moves up along n /11/, levels with $n < n_0$ are weakly excited and that is why the accepted $n_0 \sim n_{\max} \approx 10$ proved sufficient. During all the time the probability $W_{n_{\max}} = |C_{n_{\max}}|^2$ was at the level 10^{-6} , and a further decrease of n_{\min} didn't influence the dynamics of excitation. The working value of n_{\max} equal to $\approx 2n_0$ was selected in most cases. At the same time a number of control experiments with $n_{\max} \approx 2.7n_0$ was carried out. A good agreement of working and control points (see below) shows that in order to investigate electron excitation into states of the discrete spectrum within the time $\tau \approx 40$ periods of an external field we can restrict ourselves to a finite number of levels with $n_{\max} \approx 2n_0$. At values $n_{\max} \approx 2n_0$, $n_0 \approx 50$ and $\omega_0 \approx 1$ to come out into a continuous spectrum it's necessary to absorb five photons more as compared to states $n \approx n_{\max}$. That is why it is quite natural to expect that the probability of multiphoton ionization will be relatively small. Besides, the main aim of the experiments was to investigate the diffusion mechanism of excitation, occurring for discrete states only.

In the process of numerical integration probabilities $W_n = |C_n|^2$ were found, according to which the first $M_1 = \langle n - n_0 \rangle / n_0$ and the second $M_2 = \langle (n - n_0)^2 \rangle / n_0^2$ moments of distribution were determined as well as $W_{n > n_0 + 4}$ and $W_{n < n_0 - 1}$ in states with $n \in [1.5n_0]$ where brackets indicate an integer part. The step of integration Δt was determined from the condition $\epsilon \approx 2n_{\max} \Delta t \leq 0.3$ and it usually was ≈ 200 times less than an external field period. When it was reduced two times the relative change of cited values and probabilities $W_n > 10^{-4}$ during the time $\tau = 40$ was less than 1% (for $10^{-2} < W_n < 10^{-4}$ the change reached 10%). The accuracy of conservation of the complete probability $W = \sum W_n = 1$ was not worse than 0.1%. The calculation of matrix elements $Z_{nn'} = Z_{n'n}$ was done numerically according to the formulas given in /19/. At $n \approx 40$ their agreement with classical values was 2-3%. The time of integration of one period of external field at $n_0 = 30$, $n_{\max} = 63$, $\epsilon_0 = 0.04$, $\omega_0 = 1$ was ≈ 15 seconds of computer ES-1060.

Together with the quantum dynamics simulation a numerical investigation of the classical system was carried out. To exclude the peculiarity at the point $z=0$ it turned out suitable to pass on from unperturbed phase-action variables (n, λ) to new variables (n, ξ) and to a new "time" η , where the equations of motion are:

$$\begin{aligned} \frac{dn}{d\eta} &= -\epsilon n^2 \cos \omega t \sin \xi, & \frac{d\xi}{d\eta} &= 1 - \cos \xi \\ \frac{d\xi}{d\eta} &= -n^{-3} + 2\epsilon n \cos \omega t (1 - \cos \xi), & \lambda &= \xi - \sin \xi \end{aligned} \quad (5)$$

The initial distribution of classical trajectories in the phase space was a line: $n=n_0$ and equipartition distribution along the phase variable λ , which corresponded exactly to the initial conditions in the quantum case (one level with $n=n_0$). The complete number of trajectories $N=1000$. In view of the fact that in the classical system there was a finite number of levels a reflection boundary condition at the corresponding n_{\max} was introduced in the classical model. However, this had a weak effect on the values $W_{n > n_0 + 4}$ and $W_{n < n_0 - 1}$. So, for instance for $\epsilon_0 = 0.04$, $\omega_0 = 1$, $\tau = 40$ when reflection was replaced by absorption their change proved less than one standard deviation. The value of the latter was found by four groups, consisting of 250 classical trajectories equally distributed along λ . At the selected integration step the accuracy of conservation of energy in a constant field, e.i. at $\omega=0$, $t \leq 80 \pi n_0^2$, $\epsilon_0 = 0.06$ was better than 10^{-2} . It should be mentioned that the classical dynamics depends on scale variables only $\epsilon_0 = \epsilon n_0^2$, $\omega_0 = \omega n_0^2$ and at fixed values of the latter ones doesn't depend on n_0 /6,20/.

IV. The Results of the Experiments

a) One-frequency Excitation

The main group of experiments was conducted for values $n_0 = 30, 45, 66$ in the range of parameters ϵ_0, ω_0 where a strong stochasticity occurs, which leads to a diffusion excitation of a classical electron. However, despite the fact that $n_0 \gg 1$ the obtained results show that the excitation of an atom is of a fundamentally quantum nature. In fact, as one can see in Fig.1, quantum averages are close to classical ones only for a very short time $\tau < \tau^* \approx 5$ periods of an external field. When $\tau > \tau^*$ a quantum diffusion limitation is observed, because of which the excitation of a quantum atom proves much weaker than that of a classical one. We note that the given effect is not related to the finite number of selected levels.

Indeed, for $n_0=45$ when $n_{max}=n_0$ is reduced one time and a half (to $n_{max}=80$) the relative change M_2 increases from 4% when $\tau=10$ to 24% when $\tau=40$ and the ratio of the classical value M_2 to the quantum one changes from 2 to 6.




Fig. 1. Dependence of the moments of distribution on time for classical (1) and quantum (2) atoms with $\epsilon_0 = \epsilon n_0^2 = 0.04$, $\omega_0 = \omega n_0^2 = 1$, $n_0=45$, $n_{max}=97$.

Besides the effective width of distribution $\Delta n \approx 5 \ll n_{max}-n_0=52$ and the probability on levels with $n \approx n_{max}=97$ is $\sim 10^{-6}$. The classical value $(\Delta n)^2$ agrees well with the theoretical estimate (3) $(\Delta n)^2 \approx 2 \epsilon_0^2 n_0^2 \tau$. But to check it up in a more detailed way it's necessary to calculate the local coefficient D_n , that goes beyond our paper.

It should be noted, however, that the proximity of the first and the second moments of quantum distribution to their classical values doesn't at all mean that the excitation occurs due to diffusion. An example of this is the excitation from $n_0=30$ at $\epsilon_0=0.04$, $\omega_0=1$.

In this case for $\tau \leq 30$ the difference between quantum moments and classical ones is smaller than 20% (except the values $\tau=4,5,6,7$, where it comes up to 100% for M_1). Nevertheless, in the quantum case the dependence of excitation probabilities $W_{n \rightarrow n_0+k}$, $W_{n_0 \rightarrow n}$ on frequency ω_0 is of a pronounced resonance character and differs fundamentally from the classical one (Fig. 2a,c). For




Fig. 2. Dependence of probabilities of atom excitation $W_{n \rightarrow n_0+k}$ and $W_{n_0 \rightarrow n}$ on frequency ω_0 in the quantum (●) and classical (▲) case. For (a), (c) - $\epsilon_0=0.04$, $n_0=30$, $n_{max}=63$ (●)-control points for the quantum system with $n_{max}=80$; (b), (d) - $\epsilon_0=0.04$, $n_0=45$, $n_{max}=97$ (●)- $n_{max}=122$; (●)-quantum and (x)-classical values for $\epsilon_0=0.04$, $n_0=66$, $n_{max}=143$ (for $W_{n_0 \rightarrow n}$ classical values (▲) are given in Fig. 2c); the values of probabilities are taken at the moment of time $\tau = \omega t / 2 \pi = 30$. For (e), (f) - $\epsilon_0=0.06$, $n_0=45$, $\tau=12$, $n_{max}=97$ (●)-control point with $n_{max}=122$. Arrows point to the position of resonances and figures alongside to the

number of photons and the number of a resonant level. The curves are given to complete a graphic picture.

$\omega_0=0.9975$, $\tau=30$, $n > 34$ maximums of probability distribution fall on levels with $n=36, 37, 39, 41, 44, 47, 49, 52, 53, 66$. These values ($n \neq 36$) approximately correspond to unperturbed resonant transitions from the initial level $n_0=30$. For the last three values of n the necessary number of photons k is equal correspondingly to 10, 11, 12 (see Fig. 2). For $\omega_0=1.002$, $\tau=30$ the main contribution

(63%) into the probability $W_{n>n_0+4}$ comes from levels with $n=35(33\%), 36(18\%), 37(12\%)$. Despite the fact that close to the resonance $W_{n>[1.5n_0]}$ makes up about 20% of the complete probability, control experiments have shown that when $n_{max}-n_0$ increased one time and a half, the change of probabilities of excitation proves insignificant (see Fig. 2a,c).

Thus, in the investigated case the mechanism of excitation proves not a diffusion but a multiphoton one. At that up to $k=10$ photons can be effectively absorbed. For $\omega_0=0.9975, n_0=30, \tau=30, 0.03 \leq \epsilon_0 \leq 0.037$ the experimental value $k=9.973 \pm 0.006$.

For $n_0=45, \epsilon_0=0.04$ excitation also is of a pronounced resonant character (Fig. 2b,d). However, we cannot connect the observed resonances with multiphoton transitions between unperturbed levels. For $\omega_0=0.974, n > 49, \tau=30$ levels with $n=50-55, 57, 59, 63, 67, 72, 86, 93$ are the most excited. Even in resonance absolute values of probabilities of excitation happen to be much smaller than their classical values (Fig. 2). However, so far as the field ϵ_0 grows this difference decreases and at $\epsilon_0 \approx 0.06$ the probabilities become comparable (Fig. 3). Nevertheless, as is seen in Fig. 2e,f, even if $\epsilon_0=0.06$ the mechanism of excitation is of a resonant nature and results in a more effective excitation than classical diffusion does.




Fig. 3. Dependence of the probability of atom excitation on field ϵ_0 for $\omega_0=1, n_0=45, \tau=30, n_{max}=97$. Symbols (\bullet), (\ast) correspond to quantum probabilities $W_{n>[0.5n_0]}$, $W_{n>n_0+4}$ and (\blacktriangle), (\blacklozenge) to classical ones.

The dependence of the probability of excitation on the frequency of an external field proves rather strong. So, for $n_0=45, \epsilon_0=0.03, \omega_0=1, 1.3$ for $\tau=40$ the value $W_{n>[0.5n_0]} \approx 2 \cdot 10^{-3}$ and the probability of excitation is two orders of magnitude higher for $\omega_0=1/\sqrt{2} \approx 0.707, \tau=30$. In the classical system $W_{n>[1.5n_0]}$ changes in order of magnitude from 0.016 at $\omega_0=1, \tau=40$ to 0.12 at $\omega_0 \approx 0.707, \tau=30$. However, despite the fact that at $\omega_0 \approx 0.707$ the probabilities of excitation and the moments of distribution M_1 and M_2 are close to their classical values (for M_1, M_2 at $\tau \leq 30$ the difference $\leq 30\%$), the dependence $W_{n>n_0+4}, W_{n>[1.5n_0]}$ on the frequency ω_0 differs greatly from the classical one. As it is evident in Fig. 4 the excitation of levels with $n \geq 67$ occurs in a resonant way, levels with $n=69, 76, 78, 79, 80, 81, 82$ for $\omega_0 \approx 0.707$ being the most excited. All of them except $n=76$ ($W_{n=76} \approx 0.03$) have about the same probability $W_{n=76} \approx 0.014$. For the frequency




Fig. 4. Dependence of probabilities of atom excitation on the frequency ω_0 when $n_0=45, n_{max}=97$. Points (\bullet) are quantum values for $\epsilon_0=0.03, \tau=30$, (\blacktriangle)—classical values. The control point ($+$) corresponds to $n_{max}=122$. Quantum (\ast) and classical (\blacklozenge) values correspond to $\epsilon_0=0.04, \tau=15$.

$\omega_0 = 0.697$ the probabilities of excitation of these levels are one order less and do not correspond to the positions of maximums W_n . An emergence of such a group of strongly excited close states can be qualitatively explained in the following way. After a small number of periods ($\tau \approx 5$) the initial distribution spreads quickly. The width of spreading is equal $\Delta n \approx 7$, and $\langle n \rangle \approx 49$. As when neighbouring levels are nearly equidistant for all of them a simultaneous multiphoton excitation can take place. However, it should be noted that the number of photons necessary for that is rather great: $k = (E_{n_0} - E_{q,5})/\omega \approx 22$.

As the field increases up to $\epsilon_0 = 0.04$ the probability of excitation $W_{n>n_0}$ grows sharply from 0.062 to 0.26 ($\omega = 0.707$, $\tau = 15$). When the frequency changes the values $W_{n>n_0}$ are close to classical ones, but for $W_{n>n_0}$ a resonance dependence on ω_0 is still observed, and the difference from the classical value of probability is 100% (Fig. 4).

As the frequency grows, the probability of quantum system excitation becomes smaller than in the classical case. So, when $n_0 = 45$, $\epsilon_0 = 0.04$, $\omega_0 = 1.3$, $\tau = 50$ in the classical system $W_{n>n_0} = 0.359 \pm 0.013$, $W_{n>n_0} = 0.136 \pm 0.003$ and in the quantum one ($n_{max} = 97$) 0.181 and 0.051 correspondingly. Quantum moments of distribution M_1 and M_2 are 2.5 times smaller than classical ones. The same is valid for $\omega_0 = 1.25$.

The growth of the number of initially excited level up to $n_0 = 66$ doesn't lead to a qualitative change of dynamics. So, for $\epsilon_0 = 0.04$, $\omega_0 = 1.01$, $\tau = 35$ the moments of quantum distribution are rather (≈ 4 times) smaller than the classical ones. Besides, the quantum distribution breaks sharply at $n > 76$. This is its fundamental distinction from the classical one, which diminishes very slowly at great n . As a result the classical value of $W_{n>n_0}$ is about 5 times greater than the quantum one (Fig. 2d) and $W_{n>n_0} = 0.23$ is 5.4 times greater. The dependence of the probability of excitation on frequency also differs from the classical one (Fig. 2b, d). A significant role belongs here to the multiphoton mechanism of excitation. So, for $\omega_0 = 1$, $\tau = 20$ maximums of distribution W_n at $n = 60, 76, 89$ approximately correspond to $k = 7, 8, 15$ -photon transitions from initial level (Fig. 5).

b) Two-frequency Excitation

A number of experiments was carried out for the case when an external field has two harmonics: $\epsilon(t) = \epsilon(\cos \omega t + \cos \nu t)$. One frequency has been fixed $\omega n_0^2 = \omega_0 = 1$ and the second was being changed in a small range near the

values $\omega_1 = \nu n_0^2 = 0.7, 1.3$. For the former one a visible excitation of the classical system occurs at $\epsilon_0 = \epsilon n_0^2 \gg 0.015$. For smaller values spreading of the classical packet during the time $\tau = \omega t / 2\pi \leq 40$ proves insignificant (Fig. 6a). But at $\epsilon_0 = 0.02$, $\tau = 20$ there already appears a developed stochasticity, leading to a substantial excitation of the system (Fig. 6b, c). In the quantum case when ϵ grows the width of distribution grows as well (Fig. 6) which agrees qualitatively with the idea of the diffusion mechanism of excitation. The probability of excitation of levels close to n_0 ($W_{n \approx n_0}$) is close to the classical one (see Fig. 7a) and the probability of excitation of high levels $W_{n > n_0}$ is much greater than that and is of a marked resonance nature (Fig. 7c). The letter points to the fact that multiphoton excitation may also be rather substantial in the case of a wide distribution over levels (Fig. 6b). In all the investigated cases the quantum distribution turned out to be more




Fig. 5. Distribution over levels in the classical and quantum cases when $\epsilon_0 = 0.04$, $\omega_0 = 1$, $n_0 = 66$, $\tau = 20$, $n_{max} = 143$. The broken line is the quantum distribution: points correspond to cells of histogram of classical distribution with $\Delta n/n_0 = 0.05$, the smooth curve is given to complete a graphic picture.

cutting than the classical one, peaks W_n in a weak field correspond to resonance transitions (e.g. in Fig.6a the peak at $n=51$ corresponds to 5-photon resonance with $n_0=45$).

For $\omega_1 \approx 1.3$, $n_0 = 45$, $\epsilon_0 = 0.02$ the quantum system is less excited than the classical one (Fig.7b,d). So, for instance, for $\omega_1 = 1.27$, $n_0 = 45$, $\epsilon_0 = 0.02$ the classical value M_2 is 3 times less than the quantum one, and M_1 is 300 times less. When n_0 comes up to 66 probabilities of excitation are close to classical ones (see Fig.7b,d for $\omega_1 = 1.31$, $\tau = 40$), but the first moment of quantum distribution at $\tau > 10$ differs 2 or 4 times from the classical one and at $10 < \tau < 15$ has another sign. Sharp peaks




Fig.6. The same as in Fig.5 but in case of the two-frequency excitation for $n_0=45$, $\omega_0=1$, $\tau = \omega t/2\pi=20$, $n_{max}=97$. (a) $\epsilon_0 = 0.01$, $\omega_1=0.70711$; (b) $\epsilon_0 = 0.02$, $\omega_1=0.735$; (c) $\epsilon_0=0.02$, $\omega_1=0.70711$.




Fig.7. Dependence of the probability of atom excitation on frequency in case of two-frequency excitation when $n_0=45$, $\omega_0=1$, $n_{max}=97$. (a),(c)-for $\epsilon_0=0.015$ with quantum (*) and classical (◆) values; for $\epsilon_0=0.02$ correspondingly (•) and (▲), $\tau = 20$; (b),(d)- the same as (a),(c) but for $\tau = 40$. Symbols (⊙), (×) correspond to the quantum and classical systems when $n_0=66$, $\epsilon_0=0.02$, $\tau = 40$, $n_{max}=143$. For $W_n < [1.5n]$ the classical value is (▲).

($0.08 < W_n < 0.12$ for $n=61,67,69,70,71$) about 1.5-3 times exceeding the classical value W_n are observed in distribution for $\tau = 40$. So, despite the fact that in the quantum case a substantial spreading of the packet occurs (the excited number of levels $\Delta n \approx 20$, $M_2^q/M_2^{cl} \approx 0.8$) quantitative characteristics of distribution somehow differ from classical ones.

V. Discussion

Thus the carried out experiments have shown that the diffusion excitation of atom in a field of the monochromatic wave from states with $n \sim 50$ is rather suppressed. A simple estimate of values n , starting with which on the diffusion excitation will be close to the classical one, can be obtained in the following way [13]. According to (3) the number of diffusively excited levels Δn grows with the number of periods τ ($\omega \approx \Omega$) according to the law $(\Delta n)^2 \approx D_0 \tau \approx 2(\epsilon_0 n_0)^2 \tau$. At the same time Δn determines an effective number of lines in the spectrum of the wave function $N_L \sim \Delta n$. The spectrum can be considered continuous as far as $N_L \gg \tau$. As diffusion needs a continuous spectrum the time of quantum diffusion limitation is determined by the condition $N_L(\tau^*) \sim \tau^*$ which gives

$$\tau^* \sim 2(\epsilon_0 n_0)^2 \sim D_0 \quad (6)$$

At $\epsilon_0 \sim \epsilon_{cr}$ this time $\tau^* \sim n_0^2$ which agrees with the estimate given in [11]. Since $\tau \gg 1$, $\epsilon_0 \gg \epsilon_{cr}$ as well as $\epsilon_0 > 1/n_0$ (or $\epsilon > n^{-5}$) are the necessary conditions for diffusion, which corresponds to going beyond the perturbation theory and exceeding the quantum border of stability [16]. Thus, for the diffusion to go over a large time interval $\tau^* \gg 1$ in the field ϵ_0 (at $\omega_0 \approx 1$) the following values are needed

$$n_0 \gg 1/\epsilon_0 \quad (7)$$

At $\epsilon_0 \approx 0.03$ this condition is satisfied by $n_0 \gg 30$. Let's note that the number of levels in one non-linear resonance is $(\Delta n) \gg 1$ for $\epsilon \gg n^{-6}$ [11]. For $\epsilon_0 = 0.04$, $n_0 = 45$ the value $\tau^* \sim 6$ agrees well with the results of a numerical experiment (Fig. 1). But for τ^* to become comparable with the characteristic diffusion time $\tau_D \sim n_0^2/D_0 \sim 1/2\epsilon_0^2$ we need to satisfy this condition it's necessary to increase either the number of level n_0 or ϵ_0 . At $n_0 = 45$ the necessary value $\epsilon_0 \sim (2n_0)^{-4/2} \sim 0.1$. Computer experiments really point to the fact that with the growth of the field the probabilities of excitation in the classical and quantum cases become comparable. But with the growth of ϵ the probability of various multiphoton transitions grows also and can lead to a quicker atom excitation (Fig. 2e, f). A marked difference of the moments M_1, M_2 from the classical values also points to the quantum nature of excitation in the field with $\epsilon_0 = 0.06$. Thus for $n_0 = 45$, $\epsilon_0 = 0.06$, $\tau = 12$, $\omega_0 = 1$ the quantum moments are greater than the classical ones ($M_{1,2}^q/M_{1,2}^c = 1.8, 1.7$) and for $\omega_0 = 1.025$ smaller than the classical ones ($M_{1,2}^q/M_{1,2}^c$

$= 0.41, 0.56$). The given estimates belonged to the case when a one-dimensional approximation can be used, i.e. $n \gg n_L \sim 1$. If $n_{L,2} \sim n$ the number of lines in the spectrum $N_L \sim \Delta n_1 \Delta n_2 \sim (\epsilon_0 n_0)^2 \tau$. From which it is clear that the necessary condition of diffusion excitation is $\epsilon_0 \gg 1/n_0$. However when $\epsilon > \epsilon_{cr}$ the fulfilment of this condition doesn't at all denote that there will be no quantum diffusion limitation in the system (though the condition $N_L > \tau$ is fulfilled). This involves the fact that due to Kepler degeneration N_L contains groups of very close lines, corresponding to one shell, which decreases its efficient value. That is why to find the conditions of diffusion in the range $n_{1,2} \sim n$, $\epsilon > n^{-5}$, $\epsilon > \epsilon_{cr}$ (i.e. for $n \gg 50$) we will need further investigation.

When the frequency of excitation goes up the rate of diffusion slows down [11], which leads to a decrease of $\tau^* \sim D_0(\omega)/\omega$ ($n_L \ll n$). The marked growth of the probability of excitation at the decrease of frequency can be qualitatively explained in the following way. At $\epsilon_0 \approx 0.04$, $\omega_0 \approx 0.7$ the value $n = n_0$ is in resonance, the centre of which is at $n_r = n_0 \omega_0^{-4/2}$. This leads to an efficient increase of the field $\epsilon_0 \rightarrow \epsilon_0 (n_r/n_0)^2 = \epsilon_0 \omega_0^{-4/2} \approx 1.6 \epsilon_0$ and a more fast classical diffusion. This effect was observed for the classical atom in [20]. This sort of explanation is suitable for the diffusion mechanism of excitation only, since in case of the multiphoton mechanism the probability of excitation should decrease with the lowering of frequency. So, the resonance dependence on ω_0 , observed in the experiment (Fig. 4) still being obscure. A further theoretical explanation is also needed for the results on excitation from the states $n_0 = 30, 45, 66$ at $\omega_0 \approx 1$. So, for instance the theoretical formula, obtained in [21, 11] gives value $W_{n_0 \rightarrow [1.5, n_0]}$ 10 orders less than the experimental one ($n_0 = 30$, Fig. 2c).

In case of the two-frequency excitation a substantial spreading of the initial distribution occurs (Fig. 6), which corresponds qualitatively to the picture of diffusion excitation. However in this case as well a qualitative agreement with classics is observed for the probabilities $W_{n_0 \rightarrow n_0}$ only, while the dependence on frequency for $W_{n_0 \rightarrow [1.5, n_0]}$ is of a marked resonance nature (Fig. 7). Since the quantum diffusion limitation for quasiperiodical excitation shows much weaker than for the periodical one [22], the most optimum case for the diffusion mechanism is the one of two-frequency excitation. But in this case as well at $\epsilon_0 \approx 0.02$, $\omega_0, \omega_2 \sim 1$ the values $n_0 \approx 66$ are still in the quantum range (M_1 differs greatly from the classical one). The reason for this difference is evidently connected with the fact that even for $n_0 = 66$ the exceeding

of the quantum border of stability $\epsilon_0 \gg \epsilon_1 = 1/n_0$ at $\epsilon_0 = 0.02$ proves rather small. According to the results obtained for simple models /22/, it should be expected that when n_0 increases the time of quantum diffusion limitation will grow exponentially. This question, however, needs further investigation.

For one-frequency excitation quantum corrections lead to a substantial difference of distribution characteristics from the classical ones. This result testifies to the fact that the coincidence of the ionization probability of the classical system /6/ with the one obtained in the experiment /1/ is evidently accidental. In fact computer experiments have shown that for $n_0 = 66$, $\omega_0 = 0.4327$, $\epsilon_0 = 0.061$ already at $\tau = 3$ quantum moments are 1.7(M_1), 1.8(M_2) times smaller than the classical ones, and therefore the classical description of ionization is not applicable in this case. The probabilities of excitation are also smaller than the classical ones (for $\tau = 3$ the probabilities $W_{n_0, n_0} = 0.45$, $W_{n_0, n_0+1} = 0.23$ and corresponding quantum values are equal to 0.29, 0.1).

VI. Summary

The investigations carried out have shown that the excitation of atom from states with $n \leq 66$ by a monochromatic field with the frequency close to the Kepler one ($\omega \approx n^{-3}$) and amplitude $\epsilon \sim 0.04n^{-4}$ is of a fundamentally quantum character. The reason for this lies in the fact that on the one hand quantum corrections lead to a quantum diffusion limitation /12/, /13/, and on the other hand there may be strong multiphoton transitions in the atom, that result in many cases in a stronger excitation than the classical diffusion. When $\epsilon_0 \leq 0.04$, $\omega_0 \approx 1$ the probability of excitation out of the resonance may be one order of magnitude smaller than the classical one. So far as the field grows the probabilities become comparable, but differ 2-3 times from each other depending on the frequency ω_0 . A more efficient atom excitation is observed in the case when the frequency $\omega_0 < 1$. So, when $\epsilon_0 = 0.03$, $n_0 = 45$ the decrease of the frequency from $\omega_0 \approx 1$ to $\omega_0 \approx 0.7$ involves an increase of the probability W_{n_0, n_0} by 2 orders of magnitude. At an increase of the frequency there's no such effect.

In case of the two-frequency excitation in the range of stochasticity there occurs a significant spreading of the packet over unperturbed levels. But in these conditions as well when up to 20 levels are excited efficiently, the probability of excitation may have a resonant dependence from the frequency. Quantitative charac-

teristics of quantum distribution differ from the classical ones.

Thus, quantum diffusion limitation leads to the localization of the distribution over a relatively small group of unperturbed levels, from which strong multiphoton transitions start later on. The probability of the latter to be significant and cannot be explained by available theoretical estimates. The investigation of the dynamics of excitation in the region where the quantum diffusion limitation takes place can be conducted in experiments similar to /1-3/. Conditions of the one-dimensional approximation will be satisfied if the states with strongly different parabolic quantum numbers ($n_1^2 \approx n_2$) are excited. Modern experiment methods make it possible to perform this with a high accuracy /23/.

The author expresses his sincere gratitude to B.V.Chirikov, for his attention to this work and valuable comments, to N.B.Delone, F.M.Izrailev, V.P.Krainov for helpful discussions.

References

1. J.E.Bayfield, P.M.Koch, Phys. Rev. Lett. **33**, 258 (1974).
2. J.E.Bayfield, L.D.Gardner, P.M.Koch, Phys. Rev. Lett. **39**, 76 (1977).
3. J.E.Bayfield, L.D.Gardner, Y.Z.Gulkok, S.D.Sharma, Phys. Rev. A, **24**, 138 (1981).
4. L.V.Keldysh, Zh.Eksp.Teor.Fiz. **47**, 1945 (1964).
5. N.B.Delone, B.A.Zon, V.P.Krainov, Zh.Eksp.Teor.Fiz. **75**, 445 (1978).
6. J.G.Leopold, I.C.Percival, Phys. Rev. Lett. **41**, 944 (1978); J. Phys. B., **12**, 709 (1979).
7. B.I.Meerson, E.A.Oks, P.V.Sasorov, Pis'ma Zh.Eksp. Teor.Fiz. **29**, 79 (1979).
8. B.I.Meerson, Opt. i Spect. **51**, 582 (1981).
9. B.V.Chirikov, Phys. Reports, **52**, 263 (1979).
10. D.L.Shepelyansky, Opt. i Spect., **52**, 1102 (1982).
11. N.B.Delone, V.P.Krainov, D.L.Shepelyansky, Usp. Fiz. Nauk **140**, 355 (1983).
12. G.Casati, B.V.Chirikov, J.Ford, F.M.Izrailev, Lecture Notes in Physics, B., Springer **93**, 334 (1979).
13. B.V.Chirikov, F.M.Izrailev, D.L.Shepelyansky, Soviet Scientific Reviews **2C**, 209 (1981).
14. D.L.Shepelyansky, Dokl. Akad. Nauk SSSR **256**, 586 (1981).
15. S.P.Goreslavsky, N.B.Delone, V.P.Krainov, Preprint Phys. Inst. Acad. Sci. N33, Moscow, 1982.
16. E.V.Shyryak, Zh.Eksp.Teor.Fiz. **71**, 2039 (1976).

17. V.S.Edel'man, Usp. Fiz. Nauk 130, 675 (1980).
18. R.V.Jensen, Phys. Rev. Lett. 49, 1365 (1982).
19. L.D.Landau, E.M.Lifschitz, Quantum Mechanics, Moscow, "Nauka" 1974, p.750.
20. D.A.Jones, J.G.Leopold, I.C.Percival, J. Phys. B, 13, 31 (1980).
21. N.B.Delone, M.Yu.Ivanov, V.P.Krainov, Preprint Phys. Inst. Acad. Sci. N42, Moscow, 1983.
22. D.L.Sherelyansky, Physica 8D, 208 (1983).
23. P.M.Koch, D.R.Mariani, Phys. Rev. Lett. 46, 1275 (1981).

Д.Л.Шепелянский

КВАНТОВОЕ ОГРАНИЧЕНИЕ ДИФФУЗИИ ПРИ ВОЗБУЖДЕНИИ
РИДБЕРГОВСКОГО АТОМА В ПЕРЕМЕННОМ ПОЛЕ

Препринт
№ 83-61

Работа поступила - 12 мая 1983 г.

Ответственный за выпуск - С.Г.Попов
Подписано к печати 2.06-1983 г. МН 17567
Формат бумаги 60x90 Усл.2,1 печ.л., 1,7 учетно-изд.л.
Тираж 290 экз. Бесплатно. Заказ № 61.

Ротапринт ИЯЭ СО АН СССР, г.Новосибирск, 90