

Microwave ionization of hydrogen atoms

Recommend this on Google

◆ Dima Shepelyansky, Laboratoire de Physique Théorique, CNRS, Université Paul Sabatier, Toulouse

Microwave ionization of hydrogen atoms is a process of electron ionization of excited hydrogen atoms by an electromagnetic microwave field when tens or hundreds of photons are required to ionize one electron. Even if a microwave field is relatively weak this multiphoton ionization is much more efficient than a direct one-photon ionization at high photon energies (see Fig.1). Such a rapid ionization happens due to a diffusive growth of electron energy generated by dynamical chaos in the classical system. Quantum effects can suppress this diffusion with emergence of photonic localization which is similar to the Anderson localization in disordered solid state systems. The diffusive photoeffect was first observed in experiments of Bayfield and Koch (1974) [1], which happened to be first experiments performed in a regime of quantum chaos. The quantum effects of photonic localization were first observed by the group of Koch (1988) [2].

System description

The evolution of the classical system is governed by the Hamiltonian

$$H(\mathbf{p}, \mathbf{r}) = \mathbf{p}^2/2 - 1/|\mathbf{r}| - \epsilon r \cos \omega t,$$

written in canonical momentum-coordinate variables in atomic units, with ϵ , ω being amplitude and frequency of microwave field. The quantum evolution is described by the corresponding Schrödinger equation. The classical dynamics depends only on rescaled variables $\epsilon_0 = \epsilon/n_0^4$, $\omega_b = \omega/n_0^3$, and n_0 is a principal quantum number of initially excited level. For ionization an electron should absorb $N_I = n_0/2\omega_b$ photons. For typical experimental conditions with a microwave field of 10 GHz and $n_0 = 66$ [1], $\omega_b \approx 0.43$ so that about 76 photons are required to ionize one atom. A strong ionization takes place at relatively weak field amplitude being several times smaller than the static field ionization threshold $\epsilon_0 = 0.13$.

Kepler map




Figure 1: Fraction of ionized hydrogen atoms as a function of microwave frequency ω_b , measured in units of level spacing for initially excited level $n_0 = 66$; data are obtained by numerical simulations of classical (circles) and quantum (crosses) evolution; one-photon ionization threshold is marked by $\omega_b \phi$ (from [3])

The classical dynamics of one-dimensional atom is well described by the Kepler map

$$\bar{N} = N + k \sin \phi, \quad \bar{\phi} = \phi + 2\pi\omega(-2\omega\bar{N})^{-3/2},$$

where $N = -1/2\omega n^2$ is a number of photons at a given electron energy, ϕ is the phase of microwave field at the perihelion of electron orbit, bars mark new values of canonical conjugate variables after one orbital period, and $k = 2.58\epsilon/\omega^{5/3}$ is dimensionless kick amplitude, corresponding to an energy change generated by a microwave field on one electron period. This symplectic map is valid for $\omega_b \geq 1$. The second equation can be locally linearized giving $\bar{\phi} = \phi + TN$ with $T = 6\pi\omega^2 n_0^5$. This local map is equivalent to the Chirikov standard map with the chaos parameter $K = kT = \epsilon_0/\epsilon_c$. An example of the Poincare section of the phase space is shown in Fig.2, here the electron energy $E_0 = \omega N n_0^2$, expressed via the number of photons N , is shown as the phase of microwave field $\phi = \omega t$ at the moment when electron passes via the perihelion. Locally, the dynamics in the phase space can be described by the Chirikov standard map with the global chaos and diffusion appearing at $K = \epsilon_0/\epsilon_c > 1$. As a result the chaotic diffusive ionization takes place above the chaos border [4],[5],[6]




Figure 2: Poincare section of the Kepler map at $\epsilon_0 = 0.04$, $\omega_b = 3$ with rescaled electron energy $E_0 = \omega N n_0^2$ and microwave phase taken at perihelion.

$$\epsilon_0 > \epsilon_c \approx 1/(49\omega_0^{1/3}).$$

Above the chaos border the classical chaos leads to a rapid diffusive ionization which rate is significantly larger than a direct one-photon ionization, that is clearly seen in Fig.1. This diffusion process gives a sequence of absorption and emission of photons which is changes by $k \sin \phi$ at each orbital period of electron.

Chirikov localization of photonic transitions

The photonic transitions are characterized by the classical diffusion with the diffusion coefficient $D \approx k^2/2$. Under certain conditions the quantum interference effects lead to the Chirikov (or dynamical) localization of this diffusion. As for the quantum Chirikov standard map, the localization length counted in the number of photons is given by the diffusion rate [5],[6],[7]

$$l_\phi \approx D \approx k^2/2 = 3.33\epsilon^2/\omega^{10/3}.$$

The localization length can be also expressed in another form useful for a generic system with density levels ρ excited by a monochromatic field of strength ϵ and one-photon transition matrix element μ :

$$l_\phi \approx 2\pi^2 \epsilon^2 \mu^2 \rho^2.$$

The above expressions are valid for $\rho\omega > 1, l_\phi > 1$.

For the case of hydrogen atom we have

$\rho = n^3, \mu = 0.41/(\omega^{5/3} n^3)$. This localization is similar to the Anderson localization in disordered quasi-one-dimensional systems with the important difference that here we have a purely dynamical system without any disorder. Due to localization the excitation probability on high levels drops exponentially $f_N = \langle |\psi|^2 \rangle \propto \exp(-2|N - N_0|/l_\phi)$ as it is well seen for the results of numerical simulations shown in Fig.3. The excitation process is approximately described by the quantum Kepler map

$$\bar{\psi} = e^{-i\hat{H}_0} \hat{P} e^{-ik \cos \phi} \psi,$$

where $\hat{H}_0 = 2\pi[-2\omega(N_0 + \hat{N}_\phi)]^{-1/2}$, $N_0 = -1/(\omega n_0^2)$, $[N_\phi, \phi] = -i$ and \hat{P} is the projection operator on coupled states [6].




Figure 3: The Chirikov localization of photonic transitions: exponential probability distribution as a function of the number of absorbed photons $N_\phi = N_l - 1/(2\omega n^2)$ for $n_0 = 100, \epsilon_0 = 0.04, \omega_b = 3$, open circles give probability in one-photon interval, full circles are obtained from the quantum Kepler map, the straight line shows exponential localization.

The quantum delocalization border takes place when the localization length becomes larger than the number of photons required to ionize one electron $l_\phi > N_l$ that gives

$$\epsilon_0 > \epsilon_q \approx \omega_0^{7/6} / (6.6n_0)^{1/2} \approx 0.4\omega^{1/6} \omega_0.$$

At fixed microwave frequency the delocalization border is growing with an increase of initially excited level since $\epsilon_q \propto \omega_0 \propto n_0^3$. This theoretical prediction had been observed in the experiments of Koch group [2]. The experimental results are in good agreement with the numerical simulations based on the quantum Kepler map as it is shown in Fig.4 from [8]. Further experiments of Bayfield group with hydrogen atoms [9] and Walther group with Rydberg atoms [10] also confirmed the predictions of photonic localization theory.

Ionization of three-dimensional atoms

The above theory is based on a one-dimensional approximation of excited states of the hydrogen atom. In fact, as it is explained in [6], this approximation describes well the ionization process of real three-dimensional atoms if initial orbital number $l < (3/\omega)^{1/3}$. The physical origin of the Kepler map validity is related to the Kepler degeneracy of hydrogen atom due to which the dynamics in orbital momenta and conjugated phases is adiabatically slow and does not give significant influence on rapid dynamics in N, ϕ variables.

Chirikov localization and Anderson localization

The phenomenon of Anderson localization (1958) appears in disordered solids when a diffusive spreading in space, existing for classical trajectories, becomes exponentially localized due to quantum interference effects (a detailed description can be found in internal references and recommended reading). In systems of dynamical chaos there is no disorder but the classical diffusive spreading appears due to chaos. In a way similar to the Anderson localization this chaotic diffusion can be localized by quantum interference effects. This localization of dynamical quantum chaos is known in a literature as dynamical or Chirikov localization. This term stresses the dynamical origin of this phenomenon emerging in absence of any disorder. Here we discuss the appearance of Chirikov localization for a microwave ionization of hydrogen atoms. More examples of this phenomenon is given in the internal references below.

Stabilization in strong fields




Figure 4: Scaled 10% ionization threshold field ϵ_0 versus scaled microwave frequency $\omega_0 = \omega n_0^3$: experimental results of Koch group, taken from [2], are shown by circles, results of numerical simulations with the quantum Kepler map are shown by full circles, here $\omega/2\pi = 36.02 \text{ GHz}$ and initial level changes from 45 to 80, dashed curve shows quantum delocalization border for experiment conditions, dotted curve shows classical chaos border, no fit parameters (from [8]).

At very strong fields the classical dynamics of three-dimensional atom becomes stable and ionization is suppressed. This stabilization regime exists for the fields in the range $10\omega/(|m|+1) < \epsilon < 20\omega^2 n_0^2/(|m|+1)^2$, where m is initial magnetic quantum number [11]. The analogies between this stabilization, the Kapitza pendulum and channeling of charged particle beams in crystals are discussed in [11]. The stabilization theory still has not been verified experimentally.

Related physical systems

- ◆ **Ionization of chaotic Rydberg atoms:** A hydrogen atom is an integrable system and a microwave field should be relatively strong to induce chaotic diffusive ionization. However, it is possible to have atoms which are chaotic in absence of microwave field: it can be hydrogen or Rydberg atoms in a magnetic field, or Rydberg atoms in a static electric field. For such atoms a diffusive excitation can take place even when the microwave frequency is significantly smaller than the Kepler frequency so that hundreds of photons are required to ionize one electron. The properties of photonic localization for such atoms are analyzed in [12]. A similar type of photonic localization appears for microwave excitation of noninteracting electrons in metallic quantum dots of micron size, which can be viewed as artificial Rydberg atoms, the theory of photonic localization in such systems is described in [13].
- ◆ **Chaotic autoionization of molecular Rydberg states:** For Rydberg states of a molecule there is a coupling between rotations of dipole moment of charged molecular core and electron excited states. This coupling can be described in the frame of the Kepler map which under certain conditions leads to a diffusive autoionization of molecular Rydberg states as described in [14].
- ◆ **Capture of dark matter by the Solar System:** The capture of dark matter particles scattering on the Sun with a rotating planet is a process inverse to ionization. This capture process is also described by a simple map which is similar to the Kepler map. The energies of particles which can be captured and the capture cross-section are analytically determined in [15]. According to these results the cross-section diverges as an inverse particle energy being much larger than the planetary orbit area. The properties of dark matter chaos in the Solar System are analyzed in [16].

Historical notes

The striking experiments of Bayfield and Koch [1] done at Yale in 1974 remained for a long time as a theoretical puzzle. In 1978 Delone, Zon and Krainov [17] proposed to describe the excitation process by a diffusion equation in energy using random phase approximation and assuming that this approach is valid at $\epsilon > 1/n^5$, when the field perturbation becomes larger than the level spacing. This gave a correct estimate of the diffusion rate and ionization time scale but a wrong ionization border which goes to zero in the semiclassical limit. Independently, in 1978 Leopold and Percival [18] performed numerical simulations of classical dynamics showing that such an approach gives a fraction of ionized atoms being close to the experimental values. However, the discussion of dynamical chaos as the origin of strong ionization appears only in the work of Meerson, Oks and Sasorov in 1979 [19], where they give a correct estimate of the ionization threshold on the basis of the Chirikov criterion of resonance overlap for the case of $\omega_b \sim 1$. The dependence of the chaos border on frequency was found in [4]. The first signatures of quantum suppression of classical diffusive excitations have been found in [20] and later in advanced studies [21]. The numerical codes for quantum evolution developed in [20] were used in [3],[5],[6],[9],[21],[22]. The analytical theory of photonic localization was developed in

[5],[6],[21],[22] and it was confirmed by extensive numerical simulations performed in these works. The first experimental confirmations of this theory have been obtained by the Koch group [2] and later by the groups of Bayfield [9] and Walther [10]. The description of ionization by the classical and quantum Kepler map was developed in [6],[22]. The process of microwave ionization was studied also by Jensen [23], Blumel and Smilansky [24]. The classical Kepler map for the hydrogen ionization was also obtained in [25], for comet dynamics in the Solar system this map was derived by Petrosky [26]; map description of the dynamics of the comet Halley was developed by Chirikov and Vecheslavov [27]. Additional references and results for microwave ionization of excited atoms can be find in the reviews [28],[29],[30]. More recent experiments on microwave ionization of Rydberg atoms are presented in [31]. The term Chirikov localization was introduced in [32] to honor the pioneering contribution of Boris Chirikov in the discovery and investigation of this phenomenon.

References

- ◆ [1] Bayfield, J.E. and Koch, P.M. (1974). Multiphoton ionization of highly excited hydrogen atoms, *Phys. Rev. Lett.* 33: 258.
- ◆ [2] Galvez, E.J.; Sauer, B.E.; Moorman, L.; Koch, P.M. and Richards, D. (1988). Microwave ionization of H atoms: breakdown of classical dynamics for high frequencies, *Phys. Rev. Lett.* 61: 2011.
- ◆ [3] Casati, G.; Chirikov, B.V.; Guarneri, I. and Shepelyansky, D.L. (1986). New photoelectric ionization peak in the hydrogen atom, *Phys. Rev. Lett.* 57: 823.
- ◆ [4] Delone, N.B.; Krainov, V.P. and Shepelyansky, D.L. (1983). Highly excited atom in electromagnetic field, *Sov. Phys. Uspekhy* 26: 551.
- ◆ [5] Casati, G.; Chirikov, B.V.; Shepelyansky, D.L. and Guarneri, I. (1987). Relevance of classical chaos in quantum mechanics: the hydrogen atom in a monochromatic field, *Phys. Rep.* 154: 77.
- ◆ [6] Casati, G.; Guarneri, I. and Shepelyansky, D.L. (1988). Hydrogen atom in monochromatic field: chaos and dynamical photonic localization, *IEEE J. Quant. Elect.* 24: 1420.
- ◆ [7] Shepelyansky, D.L. (1987). Localization of diffusive excitation in multi-level systems, *Physica D* 28: 103.
- ◆ [8] Casati, G.; Guarneri, I. and Shepelyansky, D.L. (1990). Classical chaos, quantum localization and fluctuations: a unified view, *Physica A* 163: 205.
- ◆ [9] Bayfield, J.E.; Casati, G.; Guarneri, I. and Sokol, D.W. (1989). Localization of classically chaotic diffusion for hydrogen atoms in microwave fields, *Phys. Rev. Lett.* 63: 364.
- ◆ [10] Arndt, M.; Buchleitner, A.; Mantegna, R.N. and Walther, H. (1991). Experimental study of quantum and classical limits in microwave ionization of rubidium Rydberg atoms, *Phys. Rev. Lett.* 67: 2435.
- ◆ [11] Shepelyansky, D.L. (1994). Kramers map approach for stabilization of hydrogen atom in a monochromatic field, *Phys. Rev. A* 28: 575.
- ◆ [12] Benenti, G.; Casati, G. and Shepelyansky, D.L. (1999). Chaotic enhancement in microwave ionization of Rydberg atoms, *Eur. Phys. J. D* 5: 311.
- ◆ [13] Prosen, T. and Shepelyansky, D.L. (2005). Microwave control of transport through a chaotic mesoscopic dot, *Eur. Phys. J. B* 46: 515.
- ◆ [14] Benvenuto, F.; Casati, G. and Shepelyansky, D.L. (1994). Chaotic autoionization of molecular Rydberg states, *Phys. Rev. Lett.* 72: 1818.
- ◆ [15] Khriplovich, I.B. and Shepelyansky, D.L. (2009). Capture of dark matter by the Solar System, *Int.*

J. Mod. Phys. D 18(12): 1903.

- ◆ [16] Lages, J. and Shepelyansky, D.L. (2013). Dark matter chaos in the Solar System, *Mon. Not. Royal Astr. Soc. Lett.* 430: L25.
- ◆ [17] Delone, N.B.; Zon, B.A. and Krainov, V.P. (1978). Diffusion mechanism of ionization of highly excited atoms in an alternating electromagnetic field (*Zh. Eksp. Teor. Fiz.* 75:445 (1978)), *Sov. Phys. JETP* 48: 223.
- ◆ [18] Leopold, J.G. and Percival, I.C. (1978). Microwave Ionization and excitation of Rydberg atoms, *Phys. Rev. Lett.* 41: 944.
- ◆ [19] Meerson, B.I.; Oks, E.A. and Sasorov, P.V. (1979). Stochastic instability of an oscillator and the ionization of highly-excited atoms under the action of electromagnetic radiation (*Pis'ma Zh. Eksp. Teor. Fiz.* 29:79 (1979)), *Sov. JETP Lett.* 29: 72.
- ◆ [20] Shepelyansky, D.L. (1983). Quantum diffusion limitation at excitation of Rydberg atom in variable field, (also at Proc. Int. Conf. on Quantum Chaos (Como 1983), Plenum p.187 (1985)) *Preprint Inst. Nuclear Physics 83-61, Novosibirsk* : .
- ◆ [21] Casati, G.; Chirikov, B.V. and Shepelyansky, D.L. (1984). Quantum limitations for chaotic excitation of hydrogen atom in monochromatic field, *Phys. Rev. Lett.* 53: 2525.
- ◆ [22] Casati, G.; Guarneri, I. and Shepelyansky, D.L. (1987). Exponential photonic localization for hydrogen atom in a monochromatic field, *Phys. Rev. A* 36: 3501.
- ◆ [23] Jensen, R.V. (1982). Stochastic ionization of surface-State electrons, *Phys. Rev. Lett.* 49: 1365.
- ◆ [24] Blumel, R. and Smilansky, U. (1989). Ionization of excited hydrogen atoms by microwave fields: a test case for quantum chaos, *Phys. Scripta* 40: 386.
- ◆ [25] Gontis, V. and Kaulakys, B. (1987). Stochastic dynamics of hydrogenic atoms in the microwave field: modelling by maps and quantum description, *J. Phys. B: At. Mol. Opt. Phys.* 20: 5051.
- ◆ [26] Petrosky, T.Y. (1986). Chaos and cometary clouds in the Solar System, *Phys. Lett. A* 117: 328.
- ◆ [27] Chirikov, B.V. and Vecheslavov, V.V. (1989). Chaotic dynamics of comet Halley, *Astron. Astrophys.* 221: 146.
- ◆ [28] Jensen, R.V.; Susskind, S.M. and Sanders, M.M. (1991). Chaotic ionization of highly excited hydrogen atoms: Comparison of classical and quantum theory with experiment, *Phys. Rep.* 201: 1.
- ◆ [29] Koch, P.M. and van Leeuwen, K.H.A. (1995). The importance of resonances in microwave "ionization" of excited hydrogen atoms, *Phys. Rep.* 256: 289.
- ◆ [30] Buchleitner, A.; Delande, D. and Zakrewski, J. (2002). Non-dispersive wave packets in periodically driven quantum systems, *Phys. Rep.* 368: 409.
- ◆ [31] Gurian, J.H.; Overstreet, K.R.; Maeda, H. and Gallagher, T.F. (2010). Connecting field ionization to photoionization via 17- and 36-GHz microwave fields *Phys. Rev. A* 82: 043415.
- ◆ [32] Frahm, K.M. and Shepelyansky, D.L. (2009). Diffusion and localization for the Chirikov typical map, *Phys. Rev. B* 80: 016210.

Internal references

- ◆ Boris Chirikov and Dima Shepelyansky (2008) Chirikov standard map, Scholarpedia, 3(3):3550.
- ◆ Shmuel Fishman (2010) Anderson localization and quantum chaos maps, Scholarpedia, 5(8):9816.
- ◆ Dima Shepelyansky (2009) Chirikov criterion, Scholarpedia, 4(9):8567.
- ◆ Mark Raizen and Daniel A. Steck (2011) Cold atom experiments in quantum chaos, Scholarpedia 6(11):10468.

Recommended reading

A.J.Lichtenberg, M.A.Lieberman, "Regular and chaotic dynamics", Springer, Berlin (1992).

T.F.Gallagher, "Rydberg atoms", Cambridge University Press (1994).

L.E.Reichl, "The Transition to chaos: conservative classical systems and quantum manifestations", Springer, Berlin (2004).

Y.Imry, "Introduction to mesoscopic physics", Oxford University Press (1997)

Sponsored by: Dima Shepelyansky, Laboratoire de Physique Théorique, CNRS, Université Paul Sabatier, Toulouse

Sponsored by: Eugene M. Izhikevich, Editor-in-Chief of Scholarpedia, the peer-reviewed open-access encyclopedia

Reviewed by (http://www.scholarpedia.org/w/index.php?title=Microwave_ionization_of_hydrogen_atoms&oldid=0): Eugene M. Izhikevich, Editor-in-Chief of Scholarpedia, the peer-reviewed open-access encyclopedia

Reviewed by (http://www.scholarpedia.org/w/index.php?title=Microwave_ionization_of_hydrogen_atoms&oldid=122859): Shmuel Fishman, Department of Physics, Technion, Haifa, Israel

Accepted on: 2012-01-07 04:21:12 GMT (http://www.scholarpedia.org/w/index.php?title=Microwave_ionization_of_hydrogen_atoms&oldid=122859)

Categories: Quantum Chaos | Dynamical Systems

This page was last modified on 26 February 2013, at 15:56.



This page has been accessed 4,107 times.

Served in 1.630 secs.

"Microwave ionization of hydrogen atoms" by Dima Shepelyansky is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. Permissions beyond the scope of this license are described in the Terms of Use